

国外物理名著系列 14

The Principles of Quantum Mechanics
(4th edition)

量子力学原理 (第四版)

P. A. M. Dirac

科学出版社

北京

图字 :01-2007-2787

P. A. M. Dirac :The Principles of Quantum Mechanics

© Oxford University Press 1958

The Principles of Quantum Mechanics 4/e was originally published in English in 1958. This adaptation is published by arrangement with Oxford University Press and is for sale in the Mainland (part) of the People's Republic of China only.

量子力学原理(第四版)原书英文版于1958年出版。本改编版获得牛津大学出版社授权,仅限于在中华人民共和国大陆地区销售。版权所有,翻印必究。

图书在版编目(CIP)数据

量子力学原理:第4版=The Principles of Quantum Mechanics:4th edition;英文/(德)狄拉克(Dirac,P.A.M.)主编.—注释本.—北京:科学出版社,2008

(国外物理名著系列;14)

ISBN 978-7-03-021882-7

I. 量… II. 狄… III. 量子力学-英文 IV. 0413.1

中国版本图书馆CIP数据核字(2008)第063976号

科学出版社 出版

北京东黄城根北街16号

邮政编码:100717

<http://www.sciencep.com>

印刷

科学出版社发行 各地新华书店经销

*

2008年5月第 一 版 开本:B5(720×1000)

2008年5月第一次印刷 印张:21

印数:1~3 000 字数:385 000

定价:58.00元

(如有印装质量问题,我社负责调换〈佳倍达〉)

《国外物理名著系列》专家委员会名单

(按姓氏笔画排序)

于 涪 王鼎盛 刘正猷 刘寄星 向 涛
杨国桢 邹英华 宋菲君 张元仲 赵凯华
侯伯元 聂玉昕 阎守胜 裴寿庸 戴元本

国外物理名著系列序言

对于国内的物理学工作者和青年学生来讲，研读国外优秀的物理学著作是系统掌握物理学知识的一个重要手段。但是，在国内并不能及时、方便地买到国外的图书，且国外图书不菲的价格往往令国内的读者却步，因此，把国外的优秀物理原著引进到国内，让国内的读者能够方便地以较低的价格购买是一项意义深远的工作，将有助于国内物理学工作者和青年学生掌握国际物理学的前沿知识，进而推动我国物理学科研究和教学的发展。

为了满足国内读者对国外优秀物理学著作的需求，科学出版社启动了引进国外优秀著作的工作，出版社的这一举措得到了国内物理学界的积极响应和支持，很快成立了专家委员会，开展了选题的推荐和筛选工作，在出版社初选的书单基础上确定了第一批引进的项目，这些图书几乎涉及了近代物理学的所有领域，既有阐述学科基本理论的经典名著，也有反映某一学科专题前沿的专著。在选择图书时，专家委员会遵循了以下原则：基础理论方面的图书强调“经典”，选择了那些经得起时间检验、对物理学的发展产生重要影响、现在还不“过时”的著作（如：狄拉克的《量子力学原理》）。反映物理学某一领域进展的著作强调“前沿”和“热点”，根据国内物理学研究发展的实际情况，选择了能够体现相关学科最新进展，对有关方向的科研人员和研究生有重要参考价值的图书。这些图书都是最新版的，多数图书都是2000年以后出版的，还有相当一部分是2006年出版的新书。因此，这套丛书具有权威性、前瞻性和应用性强的特点。由于国外出版社的要求，科学出版社对部分图书进行了少量的翻译和注释（主要是目录标题和练习题），但这并不会影响图书“原汁原味”的感觉，可能还会方便国内读者的阅读和理解。

“他山之石，可以攻玉”，希望这套丛书的出版能够为国内物理学工作者和青年学生的工作和学习提供参考，也希望国内更多专家参与到这一工作中来，推荐更多的好书。



中国科学院院士
中国物理学会理事长

第四版前言

THE main change from the third edition is that the chapter on quantum electrodynamics has been rewritten. The quantum electrodynamics given in the third edition describes the motion of individual charged particles moving through the electromagnetic field, in close analogy with classical electrodynamics. It is a form of theory in which the number of charged particles is conserved and it cannot be generalized to allow of variation of the number of charged particles.

In present-day high-energy physics the creation and annihilation of charged particles is a frequent occurrence. A quantum electrodynamics which demands conservation of the number of charged particles is therefore out of touch with physical reality. So I have replaced it by a quantum electrodynamics which includes creation and annihilation of electron-positron pairs. This involves abandoning any close analogy with classical electron theory, but provides a closer description of nature. It seems that the classical concept of an electron is no longer a useful model in physics, except possibly for elementary theories that are restricted to low-energy phenomena.

P. A. M. D.

ST. JOHN'S COLLEGE, CAMBRIDGE

11 *May* 1957

NOTE TO THE REVISION OF THE FOURTH EDITION

THE opportunity has been taken of revising parts of Chapter XII ('Quantum electrodynamics') and of adding two new sections on interpretation and applications.

P. A. M. D.

ST. JOHN'S COLLEGE, CAMBRIDGE

26 *May* 1967

第一版前言

THE methods of progress in theoretical physics have undergone a vast change during the present century. The classical tradition has been to consider the world to be an association of observable objects (particles, fluids, fields, etc.) moving about according to definite laws of force, so that one could form a mental picture in space and time of the whole scheme. This led to a physics whose aim was to make assumptions about the mechanism and forces connecting these observable objects, to account for their behaviour in the simplest possible way. It has become increasingly evident in recent times, however, that nature works on a different plan. Her fundamental laws do not govern the world as it appears in our mental picture in any very direct way, but instead they control a substratum of which we cannot form a mental picture without introducing irrelevancies. The formulation of these laws requires the use of the mathematics of transformations. The important things in the world appear as the invariants (or more generally the nearly invariants, or quantities with simple transformation properties) of these transformations. The things we are immediately aware of are the relations of these nearly invariants to a certain frame of reference, usually one chosen so as to introduce special simplifying features which are unimportant from the point of view of general theory.

The growth of the use of transformation theory, as applied first to relativity and later to the quantum theory, is the essence of the new method in theoretical physics. Further progress lies in the direction of making our equations invariant under wider and still wider transformations. This state of affairs is very satisfactory from a philosophical point of view, as implying an increasing recognition of the part played by the observer in himself introducing the regularities that appear in his observations, and a lack of arbitrariness in the ways of nature, but it makes things less easy for the learner of physics. The new theories, if one looks apart from their mathematical setting, are built up from physical concepts which cannot be explained in terms of things previously known to the student, which cannot even be explained adequately in words at all. Like the fundamental concepts (e.g. proximity, identity) which every one must learn on his

arrival into the world, the newer concepts of physics can be mastered only by long familiarity with their properties and uses.

From the mathematical side the approach to the new theories presents no difficulties, as the mathematics required (at any rate that which is required for the development of physics up to the present) is not essentially different from what has been current for a considerable time. Mathematics is the tool specially suited for dealing with abstract concepts of any kind and there is no limit to its power in this field. For this reason a book on the new physics, if not purely descriptive of experimental work, must be essentially mathematical. All the same the mathematics is only a tool and one should learn to hold the physical ideas in one's mind without reference to the mathematical form. In this book I have tried to keep the physics to the forefront, by beginning with an entirely physical chapter and in the later work examining the physical meaning underlying the formalism wherever possible. The amount of theoretical ground one has to cover before being able to solve problems of real practical value is rather large, but this circumstance is an inevitable consequence of the fundamental part played by transformation theory and is likely to become more pronounced in the theoretical physics of the future.

With regard to the mathematical form in which the theory can be presented, an author must decide at the outset between two methods. There is the symbolic method, which deals directly in an abstract way with the quantities of fundamental importance (the invariants, etc., of the transformations) and there is the method of coordinates or representations, which deals with sets of numbers corresponding to these quantities. The second of these has usually been used for the presentation of quantum mechanics (in fact it has been used practically exclusively with the exception of Weyl's book *Gruppentheorie und Quantenmechanik*). It is known under one or other of the two names 'Wave Mechanics' and 'Matrix Mechanics' according to which physical things receive emphasis in the treatment, the states of a system or its dynamical variables. It has the advantage that the kind of mathematics required is more familiar to the average student, and also it is the historical method.

The symbolic method, however, seems to go more deeply into the nature of things. It enables one to express the physical laws in a neat and concise way, and will probably be increasingly used in the future as it becomes better understood and its own special mathematics gets

developed. For this reason I have chosen the symbolic method, introducing the representatives later merely as an aid to practical calculation. This has necessitated a complete break from the historical line of development, but this break is an advantage through enabling the approach to the new ideas to be made as direct as possible.

P. A. M. D.

ST. JOHN'S COLLEGE, CAMBRIDGE

29 May 1930

目 录

第 1 章 叠加原理	1
§ 1 量子理论的需要	1
§ 2 光子的极化	4
§ 3 光子的干涉	7
§ 4 叠加与不确定性.....	10
§ 5 叠加原理的数学表达.....	14
§ 6 左矢量与右矢量.....	18
第 2 章 动力学变量与可观察量	23
§ 7 线性算符.....	23
§ 8 共轭关系.....	26
§ 9 本征值与本征矢量.....	29
§ 10 可观察量	34
§ 11 可观察量的函数	41
§ 12 普遍的物理解释	45
§ 13 对易性与相容性	49
第 3 章 表象理论	53
§ 14 基矢量	53
§ 15 δ 函数	58
§ 16 基矢量的性质	62
§ 17 线性算符的表象	67
§ 18 概率幅	72
§ 19 关于可观察量函数的若干定理	76
§ 20 符号的发展	79
第 4 章 量子条件	84
§ 21 泊松括号	84
§ 22 薛定谔表象	89
§ 23 动量表象	94
§ 24 海森伯测不准原理	97
§ 25 位移算符	99
§ 26 么正变换.....	103

第 5 章 运动方程	108
§ 27 运动方程的薛定谔形式.....	108
§ 28 运动方程的海森伯形式.....	111
§ 29 定态.....	116
§ 30 自由粒子.....	118
§ 31 波包的运动.....	121
§ 32 作用量原理.....	125
§ 33 吉布斯系综.....	130
第 6 章 初等应用	136
§ 34 谐振子.....	136
§ 35 角动量.....	140
§ 36 角动量的性质.....	144
§ 37 电子的自旋.....	149
§ 38 在有心力场中的运动.....	152
§ 39 氢原子的能级.....	156
§ 40 选择定则.....	159
§ 41 氢原子的塞曼效应.....	165
第 7 章 微扰理论	167
§ 42 概述.....	167
§ 43 微扰引起的能级变化.....	168
§ 44 引起跃迁的微扰.....	172
§ 45 对辐射的应用.....	175
§ 46 与时间无关的微扰引起的跃迁.....	178
§ 47 反常塞曼效应.....	181
第 8 章 碰撞问题	185
§ 48 概述.....	185
§ 49 散射系数.....	188
§ 50 动量表象中的解.....	193
§ 51 色散散射.....	199
§ 52 共振散射.....	201
§ 53 发射与吸收.....	204
第 9 章 包含许多相同粒子的系统	207
§ 54 对称态与反对称态.....	207
§ 55 排列作为动力学变量.....	211
§ 56 排列作为运动恒量.....	213

§ 57	能级的决定	216
§ 58	对电子的应用	219
第 10 章	辐射理论	225
§ 59	玻色子系集	225
§ 60	玻色子与振子之间的联系	227
§ 61	玻色子的发射与吸收	232
§ 62	对光子的应用	235
§ 63	光子与原子间的相互作用能	239
§ 64	辐射的发射、吸收与散射	244
§ 65	费米子系集	248
第 11 章	电子的相对论性理论	253
§ 66	粒子的相对论性处理	253
§ 67	电子的波方程	254
§ 68	洛伦兹变换下的不变性	258
§ 69	自由电子的运动	261
§ 70	自旋的存在	263
§ 71	过渡到极坐标变量	267
§ 72	氢原子能级的精细结构	269
§ 73	正电子理论	273
第 12 章	量子电动力学	276
§ 74	没有物质的电磁场	276
§ 75	量子条件的相对论形式	280
§ 76	一个时刻的动力学变量	283
§ 77	补充条件	287
§ 78	电子与正电子	292
§ 79	相互作用	298
§ 80	物理的变量	302
§ 81	诠释	306
§ 82	应用	310
索引		313

CONTENTS

I. THE PRINCIPLE OF SUPERPOSITION	1
1. The Need for a Quantum Theory	1
2. The Polarization of Photons	4
3. Interference of Photons	7
4. Superposition and Indeterminacy	10
5. Mathematical Formulation of the Principle	14
6. Bra and Ket Vectors	18
II. DYNAMICAL VARIABLES AND OBSERVABLES	23
7. Linear Operators	23
8. Conjugate Relations	26
9. Eigenvalues and Eigenvectors	29
10. Observables	34
11. Functions of Observables	41
12. The General Physical Interpretation	45
13. Commutability and Compatibility	49
III. REPRESENTATIONS	53
14. Basic Vectors	53
15. The δ Function	58
16. Properties of the Basic Vectors	62
17. The Representation of Linear Operators	67
18. Probability Amplitudes	72
19. Theorems about Functions of Observables	76
20. Developments in Notation	79
IV. THE QUANTUM CONDITIONS	84
21. Poisson Brackets	84
22. Schrödinger's Representation	89
23. The Momentum Representation	94
24. Heisenberg's Principle of Uncertainty	97
25. Displacement Operators	99
26. Unitary Transformations	103
V. THE EQUATIONS OF MOTION	108
27. Schrodinger's Form for the Equations of Motion	108
28. Heisenberg's Form for the Equations of Motion	111
29. Stationary States	116
30. The Free Particle	118
31. The Motion of Wave Packets	121
32. The Action Principle	125
33. The Gibbs Ensemble	130
VI. ELEMENTARY APPLICATIONS	136
34. The Harmonic Oscillator	136
35. Angular Momentum	140

36. Properties of Angular Momentum	144
37. The Spin of the Electron	149
38. Motion in a Central Field of Force	152
39. Energy-levels of the Hydrogen Atom	156
40. Selection Rules	159
41. The Zeeman Effect for the Hydrogen Atom	165
VII. PERTURBATION THEORY	167
42. General Remarks	167
43. The Change in the Energy-levels caused by a Perturbation	168
44. The Perturbation considered as causing Transitions	172
45. Application to Radiation	175
46. Transitions caused by a Perturbation Independent of the Time	178
47. The Anomalous Zeeman Effect	181
VIII. COLLISION PROBLEMS	185
48. General Remarks	185
49. The Scattering Coefficient	188
50. Solution with the Momentum Representation	193
51. Dispersive Scattering	199
52. Resonance Scattering	201
53. Emission and Absorption	204
IX. SYSTEMS CONTAINING SEVERAL SIMILAR PARTICLES	207
54. Symmetrical and Antisymmetrical States	207
55. Permutations as Dynamical Variables	211
56. Permutations as Constants of the Motion	213
57. Determination of the Energy-levels	216
58. Application to Electrons.	219
X. THEORY OF RADIATION	225
59. An Assembly of Bosons	225
60. The Connexion between Bosons and Oscillators	227
61. Emission and Absorption of Bosons	232
62. Application to Photons	235
63. The Interaction Energy between Photons and an Atom	239
64. Emission, Absorption, and Scattering of Radiation	244
65. An Assembly of Fermions	248
XI. RELATIVISTIC THEORY OF THE ELECTRON	253
66. Relativistic Treatment of a Particle	253
67. The Wave Equation for the Electron	254
68. Invariance under a Lorentz Transformation	258
69. The Motion of a Free Electron	261
70. Existence of the Spin	263
71. Transition to Polar Variables	267
72. The Fine-structure of the Energy-levels of Hydrogen	269
73. Theory of the Positron	273

CONTENTS

xv

XII.	QUANTUM ELECTRODYNAMICS	276
	74. The Electromagnetic Field in the Absence of Matter	276
	75. Relativistic Form of the Quantum Conditions	280
	76. The Dynamical Variables at one Time	283
	77. The Supplementary Conditions	287
	78. Electrons and Positrons by Themselves	292
	79. The Interaction	298
	80. The Physical Variables	302
	81. Interpretation	306
	82. Applications	310
	INDEX	313

第 1 章 叠加原理

1. 量子理论的需要

CLASSICAL mechanics has been developed continuously from the time of Newton and applied to an ever-widening range of dynamical systems, including the electromagnetic field in interaction with matter. The underlying ideas and the laws governing their application form a simple and elegant scheme, which one would be inclined to think could not be seriously modified without having all its attractive features spoilt. Nevertheless it has been found possible to set up a new scheme, called quantum mechanics, which is more suitable for the description of phenomena on the atomic scale and which is in some respects more elegant and satisfying than the classical scheme. This possibility is due to the changes which the new scheme involves being of a very profound character and not clashing with the features of the classical theory that make it so attractive, as a result of which all these features can be incorporated in the new scheme.

The necessity for a departure from classical mechanics is clearly shown by experimental results. In the first place the forces known in classical electrodynamics are inadequate for the explanation of the remarkable stability of atoms and molecules, which is necessary in order that materials may have any definite physical and chemical properties at all. The introduction of new hypothetical forces will not save the situation, since there exist general principles of classical mechanics, holding for all kinds of forces, leading to results in direct disagreement with observation. For example, if an atomic system has its equilibrium disturbed in any way and is then left alone, it will be set in oscillation and the oscillations will get impressed on the surrounding electromagnetic field, so that their frequencies may be observed with a spectroscope. Now whatever the laws of force governing the equilibrium, one would expect to be able to include the various frequencies in a scheme comprising certain fundamental frequencies and their harmonics. This is not observed to be the case. Instead, there is observed a new and unexpected connexion between the frequencies, called Ritz's Combination Law of Spectroscopy, according to which all the frequencies can be expressed as differences between certain terms,

the number of terms being much less than the number of frequencies. This law is quite unintelligible from the classical standpoint.

One might try to get over the difficulty without departing from classical mechanics by assuming each of the spectroscopically observed frequencies to be a fundamental frequency with its own degree of freedom, the laws of force being such that the harmonic vibrations do not occur. Such a theory will not do, however, even apart from the fact that it would give no explanation of the Combination Law, since it would immediately bring one into conflict with the experimental evidence on specific heats. Classical statistical mechanics enables one to establish a general connexion between the total number of degrees of freedom of an assembly of vibrating systems and its specific heat. If one assumes all the spectroscopic frequencies of an atom to correspond to different degrees of freedom, one would get a specific heat for any kind of matter very much greater than the observed value. In fact the observed specific heats at ordinary temperatures are given fairly well by a theory that takes into account merely the motion of each atom as a whole and assigns no internal motion to it at all.

This leads us to a new clash between classical mechanics and the results of experiment. There must certainly be some internal motion in an atom to account for its spectrum, but the internal degrees of freedom, for some classically inexplicable reason, do not contribute to the specific heat. A similar clash is found in connexion with the energy of oscillation of the electromagnetic field in a vacuum. Classical mechanics requires the specific heat corresponding to this energy to be infinite, but it is observed to be quite finite. A general conclusion from experimental results is that oscillations of high frequency do not contribute their classical quota to the specific heat.

As another illustration of the failure of classical mechanics we may consider the behaviour of light. We have, on the one hand, the phenomena of interference and diffraction, which can be explained only on the basis of a wave theory; on the other, phenomena such as photo-electric emission and scattering by free electrons, which show that light is composed of small particles. These particles, which are called photons, have each a definite energy and momentum, depending on the frequency of the light, and appear to have just as real an existence as electrons, or any other particles known in physics. A fraction of a photon is never observed.

Experiments have shown that this anomalous behaviour is not peculiar to light, but is quite general. All material particles have wave properties, which can be exhibited under suitable conditions. We have here a very striking and general example of the breakdown of classical mechanics—not merely an inaccuracy in its laws of motion, but *an inadequacy of its concepts to supply us with a description of atomic events.*

The necessity to depart from classical ideas when one wishes to account for the ultimate structure of matter may be seen, not only from experimentally established facts, but also from general philosophical grounds. In a classical explanation of the constitution of matter, one would assume it to be made up of a large number of small constituent parts and one would postulate laws for the behaviour of these parts, from which the laws of the matter in bulk could be deduced. This would not complete the explanation, however, since the question of the structure and stability of the constituent parts is left untouched. To go into this question, it becomes necessary to postulate that each constituent part is itself made up of smaller parts, in terms of which its behaviour is to be explained. There is clearly no end to this procedure, so that one can never arrive at the ultimate structure of matter on these lines. So long as *big* and *small* are merely relative concepts, it is no help to explain the big in terms of the small. It is therefore necessary to modify classical ideas in such a way as to give an absolute meaning to size.

At this stage it becomes important to remember that science is concerned only with observable things and that we can observe an object only by letting it interact with some outside influence. An act of observation is thus necessarily accompanied by some disturbance of the object observed. We may define an object to be big when the disturbance accompanying our observation of it may be neglected, and small when the disturbance cannot be neglected. This definition is in close agreement with the common meanings of big and small.

It is usually assumed that, by being careful, we may cut down the disturbance accompanying our observation to any desired extent. The concepts of big and small are then purely relative and refer to the gentleness of our means of observation as well as to the object being described. In order to give an absolute meaning to size, such as is required for any theory of the ultimate structure of matter, we have to assume that *there is a limit to the fineness of our powers of observation*

and the smallness of the accompanying disturbance—a limit which is inherent in the nature of things and can never be surpassed by improved technique or increased skill on the part of the observer. If the object under observation is such that the unavoidable limiting disturbance is negligible, then the object is big in the absolute sense and we may apply classical mechanics to it. If, on the other hand, the limiting disturbance is not negligible, then the object is small in the absolute sense and we require a new theory for dealing with it.

A consequence of the preceding discussion is that we must revise our ideas of causality. Causality applies only to a system which is left undisturbed. If a system is small, we cannot observe it without producing a serious disturbance and hence we cannot expect to find any causal connexion between the results of our observations. Causality will still be assumed to apply to undisturbed systems and the equations which will be set up to describe an undisturbed system will be differential equations expressing a causal connexion between conditions at one time and conditions at a later time. These equations will be in close correspondence with the equations of classical mechanics, but they will be connected only indirectly with the results of observations. There is an unavoidable indeterminacy in the calculation of observational results, the theory enabling us to calculate in general only the probability of our obtaining a particular result when we make an observation.

2. 光子的极化

The discussion in the preceding section about the limit to the gentleness with which observations can be made and the consequent indeterminacy in the results of those observations does not provide any quantitative basis for the building up of quantum mechanics. For this purpose a new set of accurate laws of nature is required. One of the most fundamental and most drastic of these is the *Principle of Superposition of States*. We shall lead up to a general formulation of this principle through a consideration of some special cases, taking first the example provided by the polarization of light.

It is known experimentally that when plane-polarized light is used for ejecting photo-electrons, there is a preferential direction for the electron emission. Thus the polarization properties of light are closely connected with its corpuscular properties and one must ascribe a polarization to the photons. One must consider, for instance, a beam

of light plane-polarized in a certain direction as consisting of photons each of which is plane-polarized in that direction and a beam of circularly polarized light as consisting of photons each circularly polarized. Every photon is in a certain *state of polarization*, as we shall say. The problem we must now consider is how to fit in these ideas with the known facts about the resolution of light into polarized components and the recombination of these components.

Let us take a definite case. Suppose we have a beam of light passing through a crystal of tourmaline, which has the property of letting through only light plane-polarized perpendicular to its optic axis. Classical electrodynamics tells us what will happen for any given polarization of the incident beam. If this beam is polarized perpendicular to the optic axis, it will all go through the crystal; if parallel to the axis, none of it will go through; while if polarized at an angle α to the axis, a fraction $\sin^2\alpha$ will go through. How are we to understand these results on a photon basis?

A beam that is plane-polarized in a certain direction is to be pictured as made up of photons each plane-polarized in that direction. This picture leads to no difficulty in the cases when our incident beam is polarized perpendicular or parallel to the optic axis. We merely have to suppose that each photon polarized perpendicular to the axis passes unhindered and unchanged through the crystal, while each photon polarized parallel to the axis is stopped and absorbed. A difficulty arises, however, in the case of the obliquely polarized incident beam. Each of the incident photons is then obliquely polarized and it is not clear what will happen to such a photon when it reaches the tourmaline.

A question about what will happen to a particular photon under certain conditions is not really very precise. To make it precise one must imagine some experiment performed having a bearing on the question and inquire what will be the result of the experiment. Only questions about the results of experiments have a real significance and it is only such questions that theoretical physics has to consider.

In our present example the obvious experiment is to use an incident beam consisting of only a single photon and to observe what appears on the back side of the crystal. According to quantum mechanics the result of this experiment will be that sometimes one will find a whole photon, of energy equal to the energy of the incident photon, on the back side and other times one will find nothing. When one

finds a whole photon, it will be polarized perpendicular to the optic axis. One will never find only a part of a photon on the back side. If one repeats the experiment a large number of times, one will find the photon on the back side in a fraction $\sin^2\alpha$ of the total number of times. Thus we may say that the photon has a probability $\sin^2\alpha$ of passing through the tourmaline and appearing on the back side polarized perpendicular to the axis and a probability $\cos^2\alpha$ of being absorbed. These values for the probabilities lead to the correct classical results for an incident beam containing a large number of photons.

In this way we preserve the individuality of the photon in all cases. We are able to do this, however, only because we abandon the determinacy of the classical theory. The result of an experiment is not determined, as it would be according to classical ideas, by the conditions under the control of the experimenter. The most that can be predicted is a set of possible results, with a probability of occurrence for each.

The foregoing discussion about the result of an experiment with a single obliquely polarized photon incident on a crystal of tourmaline answers all that can legitimately be asked about what happens to an obliquely polarized photon when it reaches the tourmaline. Questions about what decides whether the photon is to go through or not and how it changes its direction of polarization when it does go through cannot be investigated by experiment and should be regarded as outside the domain of science. Nevertheless some further description is necessary in order to correlate the results of this experiment with the results of other experiments that might be performed with photons and to fit them all into a general scheme. Such further description should be regarded, not as an attempt to answer questions outside the domain of science, but as an aid to the formulation of rules for expressing concisely the results of large numbers of experiments.

The further description provided by quantum mechanics runs as follows. It is supposed that a photon polarized obliquely to the optic axis may be regarded as being partly in the state of polarization parallel to the axis and partly in the state of polarization perpendicular to the axis. The state of oblique polarization may be considered as the result of some kind of superposition process applied to the two states of parallel and perpendicular polarization. This implies

a certain special kind of relationship between the various states of polarization, a relationship similar to that between polarized beams in classical optics, but which is now to be applied, not to beams, but to the states of polarization of one particular photon. This relationship allows any state of polarization to be resolved into, or expressed as a superposition of, any two mutually perpendicular states of polarization.

When we make the photon meet a tourmaline crystal, we are subjecting it to an observation. We are observing whether it is polarized parallel or perpendicular to the optic axis. The effect of making this observation is to force the photon entirely into the state of parallel or entirely into the state of perpendicular polarization. It has to make a sudden jump from being partly in each of these two states to being entirely in one or other of them. Which of the two states it will jump into cannot be predicted, but is governed only by probability laws. If it jumps into the parallel state it gets absorbed and if it jumps into the perpendicular state it passes through the crystal and appears on the other side preserving this state of polarization.

3. 光子的干涉

In this section we shall deal with another example of superposition. We shall again take photons, but shall be concerned with their position in space and their momentum instead of their polarization. If we are given a beam of roughly monochromatic light, then we know something about the location and momentum of the associated photons. We know that each of them is located somewhere in the region of space through which the beam is passing and has a momentum in the direction of the beam of magnitude given in terms of the frequency of the beam by Einstein's photo-electric law—momentum equals frequency multiplied by a universal constant. When we have such information about the location and momentum of a photon we shall say that it is in a definite *translational state*.

We shall discuss the description which quantum mechanics provides of the interference of photons. Let us take a definite experiment demonstrating interference. Suppose we have a beam of light which is passed through some kind of interferometer, so that it gets split up into two components and the two components are subsequently made to interfere. We may, as in the preceding section, take an incident beam consisting of only a single photon and inquire what

will happen to it as it goes through the apparatus. This will present to us the difficulty of the conflict between the wave and corpuscular theories of light in an acute form.

Corresponding to the description that we had in the case of the polarization, we must now describe the photon as going partly into each of the two components into which the incident beam is split. The photon is then, as we may say, in a translational state given by the superposition of the two translational states associated with the two components. We are thus led to a generalization of the term 'translational state' applied to a photon. For a photon to be in a definite translational state it need not be associated with one single beam of light, but may be associated with two or more beams of light which are the components into which one original beam has been split.† In the accurate mathematical theory each translational state is associated with one of the wave functions of ordinary wave optics, which wave function may describe either a single beam or two or more beams into which one original beam has been split. Translational states are thus superposable in a similar way to wave functions.

Let us consider now what happens when we determine the energy in one of the components. The result of such a determination must be either the whole photon or nothing at all. Thus the photon must change suddenly from being partly in one beam and partly in the other to being entirely in one of the beams. This sudden change is due to the disturbance in the translational state of the photon which the observation necessarily makes. It is impossible to predict in which of the two beams the photon will be found. Only the probability of either result can be calculated from the previous distribution of the photon over the two beams.

One could carry out the energy measurement without destroying the component beam by, for example, reflecting the beam from a movable mirror and observing the recoil. Our description of the photon allows us to infer that, *after* such an energy measurement, it would not be possible to bring about any interference effects between the two components. So long as the photon is partly in one beam and partly in the other, interference can occur when the two beams are superposed, but this possibility disappears when the photon is forced entirely into

† The circumstance that the superposition idea requires us to generalize our original meaning of translational states, but that no corresponding generalization was needed for the states of polarization of the preceding section, is an accidental one with no underlying theoretical significance.

one of the beams by an observation. The other beam then no longer enters into the description of the photon, so that it counts as being entirely in the one beam in the ordinary way for any experiment that may subsequently be performed on it.

On these lines quantum mechanics is able to effect a reconciliation of the wave and corpuscular properties of light. The essential point is the association of each of the translational states of a photon with one of the wave functions of ordinary wave optics. The nature of this association cannot be pictured on a basis of classical mechanics, but is something entirely new. It would be quite wrong to picture the photon and its associated wave as interacting in the way in which particles and waves can interact in classical mechanics. The association can be interpreted only statistically, the wave function giving us information about the probability of our finding the photon in any particular place when we make an observation of where it is.

Some time before the discovery of quantum mechanics people realized that the connexion between light waves and photons must be of a statistical character. What they did not clearly realize, however, was that the wave function gives information about the probability of *one* photon being in a particular place and not the probable number of photons in that place. The importance of the distinction can be made clear in the following way. Suppose we have a beam of light consisting of a large number of photons split up into two components of equal intensity. On the assumption that the intensity of a beam is connected with the probable number of photons in it, we should have half the total number of photons going into each component. If the two components are now made to interfere, we should require a photon in one component to be able to interfere with one in the other. Sometimes these two photons would have to annihilate one another and other times they would have to produce four photons. This would contradict the conservation of energy. The new theory, which connects the wave function with probabilities for one photon, gets over the difficulty by making each photon go partly into each of the two components. Each photon then interferes only with itself. Interference between two different photons never occurs.

The association of particles with waves discussed above is not restricted to the case of light, but is, according to modern theory, of universal applicability. All kinds of particles are associated with waves in this way and conversely all wave motion is associated with

particles. Thus all particles can be made to exhibit interference effects and all wave motion has its energy in the form of quanta. The reason why these general phenomena are not more obvious is on account of a law of proportionality between the mass or energy of the particles and the frequency of the waves, the coefficient being such that for waves of familiar frequencies the associated quanta are extremely small, while for particles even as light as electrons the associated wave frequency is so high that it is not easy to demonstrate interference.

4. 叠加与不确定性

The reader may possibly feel dissatisfied with the attempt in the two preceding sections to fit in the existence of photons with the classical theory of light. He may argue that a very strange idea has been introduced—the possibility of a photon being partly in each of two states of polarization, or partly in each of two separate beams—but even with the help of this strange idea no satisfying picture of the fundamental single-photon processes has been given. He may say further that this strange idea did not provide any information about experimental results for the experiments discussed, beyond what could have been obtained from an elementary consideration of photons being guided in some vague way by waves. What, then, is the use of the strange idea?

In answer to the first criticism it may be remarked that the main object of physical science is not the provision of pictures, but is the formulation of laws governing phenomena and the application of these laws to the discovery of new phenomena. If a picture exists, so much the better; but whether a picture exists or not is a matter of only secondary importance. In the case of atomic phenomena no picture can be expected to exist in the usual sense of the word 'picture', by which is meant a model functioning essentially on classical lines. One may, however, extend the meaning of the word 'picture' to include any way of looking at the fundamental laws which makes their self-consistency obvious. With this extension, one may gradually acquire a picture of atomic phenomena by becoming familiar with the laws of the quantum theory.

With regard to the second criticism, it may be remarked that for many simple experiments with light, an elementary theory of waves and photons connected in a vague statistical way would be adequate

to account for the results. In the case of such experiments quantum mechanics has no further information to give. In the great majority of experiments, however, the conditions are too complex for an elementary theory of this kind to be applicable and some more elaborate scheme, such as is provided by quantum mechanics, is then needed. The method of description that quantum mechanics gives in the more complex cases is applicable also to the simple cases and although it is then not really necessary for accounting for the experimental results, its study in these simple cases is perhaps a suitable introduction to its study in the general case.

There remains an overall criticism that one may make to the whole scheme, namely, that in departing from the determinacy of the classical theory a great complication is introduced into the description of Nature, which is a highly undesirable feature. This complication is undeniable, but it is offset by a great simplification, provided by the general *principle of superposition of states*, which we shall now go on to consider. But first it is necessary to make precise the important concept of a 'state' of a general atomic system.

Let us take any atomic system, composed of particles or bodies with specified properties (mass, moment of inertia, etc.) interacting according to specified laws of force. There will be various possible motions of the particles or bodies consistent with the laws of force. Each such motion is called a *state* of the system. According to classical ideas one could specify a state by giving numerical values to all the coordinates and velocities of the various component parts of the system at some instant of time, the whole motion being then completely determined. Now the argument of pp. 3 and 4 shows that we cannot observe a *small* system with that amount of detail which classical theory supposes. The limitation in the power of observation puts a limitation on the number of data that can be assigned to a state. Thus a state of an atomic system must be specified by fewer or more indefinite data than a complete set of numerical values for all the coordinates and velocities at some instant of time. In the case when the system is just a single photon, a state would be completely specified by a given translational state in the sense of § 3 together with a given state of polarization in the sense of § 2.

A state of a system may be defined as an undisturbed motion that is restricted by as many conditions or data as are theoretically possible without mutual interference or contradiction. In practice

the conditions could be imposed by a suitable preparation of the system, consisting perhaps in passing it through various kinds of sorting apparatus, such as slits and polarimeters, the system being left undisturbed after the preparation. The word 'state' may be used to mean either the state at one particular time (after the preparation), or the state throughout the whole of time after the preparation. To distinguish these two meanings, the latter will be called a 'state of motion' when there is liable to be ambiguity.

The general principle of superposition of quantum mechanics applies to the states, with either of the above meanings, of any one dynamical system. It requires us to assume that between these states there exist peculiar relationships such that whenever the system is definitely in one state we can consider it as being partly in each of two or more other states. The original state must be regarded as the result of a kind of *superposition* of the two or more new states, in a way that cannot be conceived on classical ideas. Any state may be considered as the result of a superposition of two or more other states, and indeed in an infinite number of ways. Conversely any two or more states may be superposed to give a new state. The procedure of expressing a state as the result of superposition of a number of other states is a mathematical procedure that is always permissible, independent of any reference to physical conditions, like the procedure of resolving a wave into Fourier components. Whether it is useful in any particular case, though, depends on the special physical conditions of the problem under consideration.

In the two preceding sections examples were given of the superposition principle applied to a system consisting of a single photon. § 2 dealt with states differing only with regard to the polarization and § 3 with states differing only with regard to the motion of the photon as a whole.

The nature of the relationships which the superposition principle requires to exist between the states of any system is of a kind that cannot be explained in terms of familiar physical concepts. One cannot in the classical sense picture a system being partly in each of two states and see the equivalence of this to the system being completely in some other state. There is an entirely new idea involved, to which one must get accustomed and in terms of which one must proceed to build up an exact mathematical theory, without having any detailed classical picture.

When a state is formed by the superposition of two other states, it will have properties that are in some vague way intermediate between those of the two original states and that approach more or less closely to those of either of them according to the greater or less 'weight' attached to this state in the superposition process. The new state is completely defined by the two original states when their relative weights in the superposition process are known, together with a certain phase difference, the exact meaning of weights and phases being provided in the general case by the mathematical theory. In the case of the polarization of a photon their meaning is that provided by classical optics, so that, for example, when two perpendicularly plane polarized states are superposed with equal weights, the new state may be circularly polarized in either direction, or linearly polarized at an angle $\frac{1}{4}\pi$, or else elliptically polarized, according to the phase difference.

The non-classical nature of the superposition process is brought out clearly if we consider the superposition of two states, A and B , such that there exists an observation which, when made on the system in state A , is certain to lead to one particular result, a say, and when made on the system in state B is certain to lead to some different result, b say. What will be the result of the observation when made on the system in the superposed state? The answer is that the result will be sometimes a and sometimes b , according to a probability law depending on the relative weights of A and B in the superposition process. It will never be different from both a and b . *The intermediate character of the state formed by superposition thus expresses itself through the probability of a particular result for an observation being intermediate between the corresponding probabilities for the original states,† not through the result itself being intermediate between the corresponding results for the original states.*

In this way we see that such a drastic departure from ordinary ideas as the assumption of superposition relationships between the states is possible only on account of the recognition of the importance of the disturbance accompanying an observation and of the consequent indeterminacy in the result of the observation. When an observation is made on any atomic system that is in a given state,

† The probability of a particular result for the state formed by superposition is not always intermediate between those for the original states in the general case when those for the original states are not zero or unity, so there are restrictions on the 'intermediateness' of a state formed by superposition.

in general the result will not be determinate, i.e., if the experiment is repeated several times under identical conditions several different results may be obtained. It is a law of nature, though, that if the experiment is repeated a large number of times, each particular result will be obtained in a definite fraction of the total number of times, so that there is a definite *probability* of its being obtained. This probability is what the theory sets out to calculate. Only in special cases when the probability for some result is unity is the result of the experiment determinate.

The assumption of superposition relationships between the states leads to a mathematical theory in which the equations that define a state are linear in the unknowns. In consequence of this, people have tried to establish analogies with systems in classical mechanics, such as vibrating strings or membranes, which are governed by linear equations and for which, therefore, a superposition principle holds. Such analogies have led to the name 'Wave Mechanics' being sometimes given to quantum mechanics. It is important to remember, however, that *the superposition that occurs in quantum mechanics is of an essentially different nature from any occurring in the classical theory*, as is shown by the fact that the quantum superposition principle demands indeterminacy in the results of observations in order to be capable of a sensible physical interpretation. The analogies are thus liable to be misleading.

5. 叠加原理的数学表达

A profound change has taken place during the present century in the opinions physicists have held on the mathematical foundations of their subject. Previously they supposed that the principles of Newtonian mechanics would provide the basis for the description of the whole of physical phenomena and that all the theoretical physicist had to do was suitably to develop and apply these principles. With the recognition that there is no logical reason why Newtonian and other classical principles should be valid outside the domains in which they have been experimentally verified has come the realization that departures from these principles are indeed necessary. Such departures find their expression through the introduction of new mathematical formalisms, new schemes of axioms and rules of manipulation, into the methods of theoretical physics.

Quantum mechanics provides a good example of the new ideas. It

requires the states of a dynamical system and the dynamical variables to be interconnected in quite strange ways that are unintelligible from the classical standpoint. The states and dynamical variables have to be represented by mathematical quantities of different natures from those ordinarily used in physics. The new scheme becomes a precise physical theory when all the axioms and rules of manipulation governing the mathematical quantities are specified and when in addition certain laws are laid down connecting physical facts with the mathematical formalism, so that from any given physical conditions equations between the mathematical quantities may be inferred and vice versa. In an application of the theory one would be given certain physical information, which one would proceed to express by equations between the mathematical quantities. One would then deduce new equations with the help of the axioms and rules of manipulation and would conclude by interpreting these new equations as physical conditions. The justification for the whole scheme depends, apart from internal consistency, on the agreement of the final results with experiment.

We shall begin to set up the scheme by dealing with the mathematical relations between the states of a dynamical system at one instant of time, which relations will come from the mathematical formulation of the principle of superposition. The superposition process is a kind of additive process and implies that states can in some way be added to give new states. The states must therefore be connected with mathematical quantities of a kind which can be added together to give other quantities of the same kind. The most obvious of such quantities are vectors. Ordinary vectors, existing in a space of a finite number of dimensions, are not sufficiently general for most of the dynamical systems in quantum mechanics. We have to make a generalization to vectors in a space of an infinite number of dimensions, and the mathematical treatment becomes complicated by questions of convergence. For the present, however, we shall deal merely with some general properties of the vectors, properties which can be deduced on the basis of a simple scheme of axioms, and questions of convergence and related topics will not be gone into until the need arises.

It is desirable to have a special name for describing the vectors which are connected with the states of a system in quantum mechanics, whether they are in a space of a finite or an infinite number of

dimensions. We shall call them *ket vectors*, or simply *kets*, and denote a general one of them by a special symbol $|\rangle$. If we want to specify a particular one of them by a label, A say, we insert it in the middle, thus $|A\rangle$. The suitability of this notation will become clear as the scheme is developed.

Ket vectors may be multiplied by complex numbers and may be added together to give other ket vectors, e.g. from two ket vectors $|A\rangle$ and $|B\rangle$ we can form

$$c_1 |A\rangle + c_2 |B\rangle = |R\rangle, \quad (1)$$

say, where c_1 and c_2 are any two complex numbers. We may also perform more general linear processes with them, such as adding an infinite sequence of them, and if we have a ket vector $|x\rangle$, depending on and labelled by a parameter x which can take on all values in a certain range, we may integrate it with respect to x , to get another ket vector

$$\int |x\rangle dx = |Q\rangle$$

say. A ket vector which is expressible linearly in terms of certain others is said to be *dependent* on them. A set of ket vectors are called *independent* if no one of them is expressible linearly in terms of the others.

We now assume that *each state of a dynamical system at a particular time corresponds to a ket vector, the correspondence being such that if a state results from the superposition of certain other states, its corresponding ket vector is expressible linearly in terms of the corresponding ket vectors of the other states, and conversely*. Thus the state R results from a superposition of the states A and B when the corresponding ket vectors are connected by (1).

The above assumption leads to certain properties of the superposition process, properties which are in fact necessary for the word 'superposition' to be appropriate. When two or more states are superposed, the order in which they occur in the superposition process is unimportant, so the superposition process is symmetrical between the states that are superposed. Again, we see from equation (1) that (excluding the case when the coefficient c_1 or c_2 is zero) if the state R can be formed by superposition of the states A and B , then the state A can be formed by superposition of B and R , and B can be formed by superposition of A and R . The superposition relationship is symmetrical between all three states A , B , and R .